arXiv:math/0405503v5 [math.NT] 7 Sep 2007

GALOIS MODULE STRUCTURE OF MILNOR K-THEORY IN CHARACTERISTIC p

GANESH BHANDARI, NICOLE LEMIRE*, JÁN MINÁČ†,
AND JOHN SWALLOW‡

Abstract. Let E be a cyclic extension of degree p^n of a field F of characteristic p. Using arithmetic invariants of E/F we determine k_mE, the Milnor K-groups K_mE modulo p, as $\mathbb{F}_p[\mathrm{Gal}(E/F)]$-modules for all $m \in \mathbb{N}$. In particular, we show that each indecomposable summand of k_mE has $\mathbb{F}_p$-dimension a power of p. That all powers p^i, $i = 0, 1, \ldots, n$, occur for suitable examples is shown in a subsequent paper [MSS2], where additionally the main result of this paper becomes an essential induction step in the determination of K_mE/p^sK_mE as $(\mathbb{Z}/p^s\mathbb{Z})[\mathrm{Gal}(E/F)]$-modules for all $m, s \in \mathbb{N}$.

In the 1960s Z. I. Borevič and D. K. Faddeev classified the possible $\mathbb{F}_p[\mathrm{Gal}(E/F)]$-modules k_1E, where E is a cyclic extension of degree p^n of a local field F. (See [Bo].) This result has recently been generalized: first in the $n = 1$ case, replacing the local field F by any field containing a primitive pth-root of unity [MS]; and later to all base fields F and all cyclic extensions E of degree p^n [MSS1]. Then three of the authors obtained a generalization from k_1 to k_m, handling all $\mathbb{F}_p[\mathrm{Gal}(E/F)]$-modules k_mE, $m \in \mathbb{N}$, provided that $n = 1$ and F contains a primitive pth-root of unity [LMS]. This result, in turn, led to new restrictions on possible absolute Galois pro-p groups [BLMS].

The particular case of fields of characteristic p is important not only for its intrinsic value but also because the case arises naturally in considering the residue fields of valuations on fields of characteristic 0. This case, however, has been treated only as far as k_1E for E/F cyclic

---

Date: September 5, 2007.

2000 Mathematics Subject Classification. 19D45, 12F10.

Key words and phrases. Milnor K-groups modulo p, cyclic extensions, Galois modules.

*Research supported in part by NSERC grant R3276A01.

†Research supported in part by NSERC grant R0370A01, and by a Distinguished Research Professorship at the University of Western Ontario.

‡Research supported in part by NSA grant MDA904-02-1-0061 and by NSF grant DMS-0600122.

of degree p^n [MSS1]. Twenty years ago Bloch, Gabber, and Kato established an isomorphism between, on one hand, Milnor K-theory modulo p in characteristic p and, on the other, the kernel of the Artin-Schreier operator defined on the exterior algebra on Kähler differentials [BK]. Not long afterward, two significant papers by Izhboldin considered Milnor K-theory in characteristic p and, using the previous work of Bloch, Gabber, and Kato, ascertained its important Galois module properties [I1, I2].

In this paper we establish that these two results of Izhboldin are enough to determine precisely the Galois module structure of the Milnor groups K_mE mod p when E is a cyclic extension of pth-power degree of a field F of characteristic p. The result is simple and useful: these modules are direct sums of trivial modules and modules free over some quotient of the Galois group. Equivalently, the dimensions over $\mathbb{F}_p$ of the indecomposable $\mathbb{F}_p[\mathrm{Gal}(E/F)]$-modules occurring as direct summands of K_mE/pK_mE are all powers of p. In [MSS2] it is shown that in fact each indecomposable summand of dimension p^i, $i = 0, 1, \dots, n$, occurs in suitable examples.

Our decomposition of the $\mathbb{F}_p[\mathrm{Gal}(E/F)]$-module k_mE into indecomposables in Theorem 2 is not canonical, although the ranks of the summands which are free over the various quotients of $\mathrm{Gal}(E/F)$ are invariants of the $\mathbb{F}_p[\mathrm{Gal}(E/F)]$-module k_mE. Still, our results do have a canonical formulation, stemming from Lemma 4, which is in fact equivalent to Theorem 2. In particular, the filtration of $(k_mE)^{\mathrm{Gal}(E/F)} \simeq k_mF$, induced by the images of successive powers of the maximal ideal of $\mathbb{F}_p[\mathrm{Gal}(E/F)]$, is determined by the images of the norm maps $k_mE_i \to k_mF$, where E_i runs through the intermediate fields between E and F.

It is possible that one could use the results in [J] to obtain alternative proofs, but we preferred a more self-contained, short derivation instead.

In [MSS2] our main result is used as a critical first induction step to determine K_mE/p^sK_mE as a $(\mathbb{Z}/p^s\mathbb{Z})[\mathrm{Gal}(E/F)]$-module for all $m, s \in \mathbb{N}$. The classification problem of $(\mathbb{Z}/p^s\mathbb{Z})G$-modules for cyclic G is nontrivial and in fact still not complete, despite some important results when, for instance, the module is free over $\mathbb{Z}/p^s\mathbb{Z}$. (See [T] and the references contained therein.) However, one can describe the $(\mathbb{Z}/p^s\mathbb{Z})[\mathrm{Gal}(E/F)]$-module K_mE/p^sK_mE completely and simply.

We assume in what follows that all fields have characteristic p and that m is a fixed natural number. For a field F, let K_mF denote the mth Milnor K-group of F and $k_mF = K_mF/pK_mF$. (See, for instance, [Ma] and [Mi, IX.1].) If E/F is a Galois extension of fields, let $G = \mathrm{Gal}(E/F)$ denote the associated Galois group. When G is a cyclic group we write $G = \langle\sigma\rangle$, with a suitable fixed generator σ. For the sake of simplicity we write ρ instead of $\sigma - 1$. We write $i_E\colon K_mF \to K_mE$ and $N_{E/F}\colon K_mE \to K_mF$ for the natural map induced by the inclusion and the norm maps, and we use the same notation for the induced maps modulo p between k_mF and k_mE. In order to avoid possible confusion, in a few instances we write $i_{F,E}$ instead of i_E.

Izhboldin's results are as follows.

Lemma 1 ([I2, Lemma 2.3])**.** *Suppose E/F is cyclic of degree p. Then $i_E\colon k_mF \to (k_mE)^G$ is an isomorphism.*

Theorem 1 (Hilbert 90 for Milnor K-theory: [I1, Corollary of Proposition 5], [I2, Theorem D])**.** *Suppose E/F is cyclic of pth-power degree. Then the following sequence is exact:*

$$K_mE \xrightarrow{1-\sigma} K_mE \xrightarrow{N_{E/F}} K_mF.$$

Now suppose E/F is cyclic of degree p^n. For $i = 0, 1, \ldots, n$, let E_i/F be the subextension of degree p^i of E/F, $H_i = \mathrm{Gal}(E/E_i)$, and $G_i = \mathrm{Gal}(E_i/F)$. Our main result is the following

Theorem 2. *There exists an isomorphism of $\mathbb{F}_pG$-modules $k_mE \simeq \oplus_{i=0}^{n} Y_i$, where*

- *Y_n is a free $\mathbb{F}_pG$-module of rank $\dim_{\mathbb{F}_p} N_{E/F}k_mE$,*
- *Y_i, $0 < i < n$, is a free $\mathbb{F}_pG_i$-module of rank*

$$\dim_{\mathbb{F}_p} N_{E_i/F}k_mE_i/N_{E_{i+1}/F}k_mE_{i+1}, \text{ and}$$

- *Y_0 is a trivial $\mathbb{F}_pG$-module of rank $\dim_{\mathbb{F}_p} k_mF/N_{E_1/F}k_mE_1$.*

Our proof of Theorem 2 relies on a filtration lemma, Lemma 4, which shows that certain elements in k_mE are expressible as norms. That the various norm groups $N_{E_i/F}k_mE_i$ contain enough elements is, in fact, the key to this lemma. In section 1 we present technical results on $\mathbb{F}_pG$-modules. In section 2 we state Lemma 4, prove that Theorem 2 is a corollary to Lemma 4, and prove basic lemmas in preparation for its proof. This proof is finally presented in section 3.

1. $\mathbb{F}_pG$-MODULES

For the reader's convenience, after introducing some notation, we recall in this section some basic elementary facts about $\mathbb{F}_pG$-modules. (See [C] and [L].)

Let G be a cyclic group of order p^n with generator σ. We denote $\sigma - 1$ as ρ. For an $\mathbb{F}_pG$-module U, let U^G denote the submodule of U fixed by G. For an arbitrary element $u \in U$, we say that the length $l(u)$ is the dimension over $\mathbb{F}_p$ of the $\mathbb{F}_pG$-submodule $\langle u \rangle$ of U generated by u. Then we have

$$\rho^{l(u)-1}\langle u \rangle = \langle u \rangle^G \neq \{0\} \quad \text{and} \quad \rho^{l(u)}\langle u \rangle = \{0\}.$$

If we want to stress the dependence of length $l(u)$ on G, we write $l_G(u)$. As usual, if U is a free $\mathbb{F}_pG$-module with $U = \oplus_{i \in \mathcal{I}} \mathbb{F}_pG$, we say that U is a module of rank $|\mathcal{I}|$. Denote by N the operator ρ^{p^n-1} acting on module U.

We use the following general lemma about $\mathbb{F}_pG$-modules. The simple proof of this lemma is omitted.

Lemma 2 (Exclusion Lemma)**.** *Let M_i, $i \in \mathcal{I}$, be a family of $\mathbb{F}_pG$-modules contained in a common $\mathbb{F}_pG$-module N. Suppose that the $\mathbb{F}_p$-vector subspace R of N generated by all M_i^G has the form $R = \oplus_{i \in \mathcal{I}} M_i^G$. Then the $\mathbb{F}_pG$-module M generated by M_i, $i \in \mathcal{I}$, has the form $M = \oplus_{i \in \mathcal{I}} M_i$.*

We also use the following result about the restriction of a cyclic $\mathbb{F}_pG$-module to a subgroup.

Lemma 3 (Restriction Lemma)**.** *Let Y be a cyclic $\mathbb{F}_pG$-module of length l generated by γ. Then $l_{H_k}(\gamma)$ is the unique integer such that $l = (l_{H_k}(\gamma) - 1)p^k + r$ for some unique integer $1 \leq r \leq p^k$. Moreover,*

$$Y \simeq V^r_{l_{H_k}(\gamma)} \oplus V^{p^k-r}_{l_{H_k}(\gamma)-1}$$

where V_i is a cyclic $\mathbb{F}_pH_k$-module of length i.

Proof. The formula for l_{H_k} follows immediately from the definition of length. Since $\{\rho^i\gamma \mid 0 \leq i \leq l-1\}$ is an $\mathbb{F}_p$-basis for $Y = \mathbb{F}_pG\gamma$, then

$$\mathbb{F}_pG\gamma\big|_{H_k} \;=\; \oplus_{i=0}^{p^k-1}\mathbb{F}_pH_k\rho^i\gamma \;\simeq\; V^r_{l_{H_k}(\gamma)} \oplus V^{p^k-r}_{l_{H_k}(\gamma)-1}$$

since $l_{H_k}(\rho^j\gamma) = l_{H_k}(\gamma)$ if $0 \leq j < r$ and $l_{H_k}(\rho^j\gamma) = l_{H_k}(\gamma) - 1$ if $r \leq j \leq p^k - 1$. □

Finally, we need a general structure proposition about $\mathbb{F}_pG$-modules that shows that the structure of an $\mathbb{F}_pG$-module X can be determined from the natural filtration on X^G obtained by taking the intersection of X^G with the images of X under the successive powers of the augmentation ideal of $\mathbb{F}_pG$. The proposition below is proved for cyclic p-groups as [LMS, Proposition 2]. The generalization to cyclic groups of order p^n is automatic.

Proposition 1. *Let X be an $\mathbb{F}_pG$-module. Set $L_{p^n} = \rho^{p^n-1}X$, and for $1 \le i < p^n$, suppose that L_i is an $\mathbb{F}_p$-complement of $\rho^i X \cap X^G$ in $\rho^{i-1}X \cap X^G$.*

Then there exist $\mathbb{F}_pG$-modules X_i, $i = 1, 2, \dots, p^n$, such that

1. $X = \bigoplus_{i=1}^{p^n} X_i$,
2. $X_i^G = L_i$ *for* $i = 1, 2, \dots, p^n$,
3. *each* X_i *is a direct sum of* $\dim_{\mathbb{F}_p}(L_i)$ *cyclic* $\mathbb{F}_pG$*-modules of length* i*, and*
4. *for each* $i = 1, 2, \dots, p^n$*, there exists an* $\mathbb{F}_p$*-submodule* Y_i *of* X_i *with* $\dim_{\mathbb{F}_p}(Y_i) = \dim_{\mathbb{F}_p}(L_i)$ *such that* $\mathbb{F}_pGY_i = X_i$.

2. Milnor k-groups

If $\alpha \in K_mE$ we write $\bar{\alpha}$ for the class of α in k_mE. For $\gamma \in K_mE$, let $l(\gamma)$ denote the dimension over $\mathbb{F}_p$ of the $\mathbb{F}_pG$-submodule $\langle\bar{\gamma}\rangle$ of k_mE generated by $\bar{\gamma}$. Because $\rho^{p^n-1} = 1 + \sigma + \cdots + \sigma^{p^n-1}$ in $\mathbb{F}_pG$, we may use $i_EN_{E/F}$ and N interchangeably on k_mE.

We will prove the following key lemma by induction on n.

Lemma 4. *Let $1 \le j \le p^n$ and let $0 \le i \le n$ be minimal such that $j \le p^i$. Then*

$$\rho^{j-1}k_mE \cap (k_mE)^G = i_EN_{E_i/F}k_mE_i.$$

Remark. Note that Lemma 1 proves this result for $j = 1$ and $n = 1$. Note also that for $j = p^n$, the result holds by definition. Moreover, for i minimal such that $j \le p^i$,

$$\begin{aligned} i_EN_{E_i/F}k_mE_i &= i_{E_i,E}i_{F,E_i}N_{E_i/F}k_mE_i = \rho^{p^i-1}(i_{E_i,E}k_mE_i) \\ &\subseteq \rho^{p^i-1}k_mE \cap (k_mE)^G \subseteq \rho^{j-1}k_mE \cap (k_mE)^G, \end{aligned}$$

so it suffices to show the inclusion in the opposite direction.

Before proving this lemma, we show

Reduction to Lemma 4. Theorem 2 holds for those cyclic extensions of degree p^n for which Lemma 4 holds.

Proof. We apply Proposition 1 to k_mE. Set $L_{p^n} = i_E N_{E/F} k_mE$, and for $0 \leq i < n$, define L_{p^i} as an $\mathbb{F}_p$-complement of $i_E N_{E_{i+1}/F} k_m E_{i+1}$ in $i_E N_{E_i/F} k_m E_i$. We also set $L_k = \{0\}$ for each $k \in \{1, 2, \dots, p^n\}$ which is not a power of p. Note that $(k_mE)^G = i_E k_m F$ by Lemma 4 applied to $j = 1$. Then, Lemma 4 shows that the $\mathbb{F}_p$-modules L_j, $j = 1, \dots, p^n$, satisfy the conditions of the proposition. By the proposition we have a decomposition of k_mE into a direct sum of $\mathbb{F}_pG$-modules

$$k_mE = \oplus_{i=0}^n Y_i,$$

where $Y_i^G = L_{p^i}$ and Y_i is a direct sum of

$$\begin{aligned}\dim_{\mathbb{F}_p} L_{p^i} &= \dim_{\mathbb{F}_p} i_E N_{E_i/F} k_m E_i / i_E N_{E_{i+1}/F} k_m E_{i+1} \\ &= \dim_{\mathbb{F}_p} N_{E_i/F} k_m E_i / N_{E_{i+1}/F} k_m E_{i+1}\end{aligned}$$

cyclic $\mathbb{F}_pG$-modules of length p^i, using Lemma 1 for the last equality. Equivalently, $Y_i \simeq \oplus_{T_i} \mathbb{F}_p(G/H_i)$, where the cardinal number of T_i is $\dim_{\mathbb{F}_p} L_{p^i}$, and therefore Y_i is a free $\mathbb{F}_pG_i$-module of the same rank. □

We now prove several lemmas as partial results toward the proof of Lemma 4. Note that it is only in these lemmas that Theorem 1 is used in this paper.

Lemma 5. *Lemma 4 holds for all cyclic extensions E/F of degree $p > 2$.*

Proof. Let $G = \mathrm{Gal}(E/F)$. By the remark to Lemma 4, it suffices to show the left-to-right inclusion of Lemma 4. Let $\gamma \in K_mE$ such that $0 \neq \rho^{l-1}\bar{\gamma} \in \rho^{l-1} k_mE \cap (k_mE)^G$ where $1 < l \leq p$. Then $l(\gamma) = l$. We show by induction on i, $l \leq i \leq p$, that there exists $\alpha_i \in K_mE$ such that

$$\langle \rho^{i-1}\bar{\alpha}_i \rangle = \langle \bar{\gamma} \rangle^G.$$

Then since $\rho^{p-1}\bar{\alpha}_p = i_E N_{E/F} \bar{\alpha}_p$,the proof will be complete. If $i = l$ then $\alpha_i = \gamma$ suffices. Assume now that $l \leq i < p$ and our statement is true for i.

Set $c := N_{E/F}\alpha_i$. Since $i_E\bar{c} = N\bar{\alpha}_i = \rho^{p-1}\bar{\alpha}_i$ and $i < p$, we conclude that $i_E\bar{c} = 0$. By the injectivity of i_E from Lemma 1, $\bar{c} = 0$. Therefore there exists $f \in K_mF$ such that $c = pf$ in K_mF.

We calculate

$$N_{E/F}(\alpha_i - i_E(f)) = c - pf = 0.$$

By Theorem 1, there exists $\omega \in K_mE$ such that $\rho\omega = \alpha_i - i_E(f)$. Hence $\rho^2\omega = \rho\alpha_i$. Since $i \geq 2$,

$$\langle \rho^i \bar{\omega} \rangle = \langle \rho^{i-1}\bar{\alpha}_i \rangle = \langle \bar{\gamma} \rangle^G$$

and we can set $\alpha_{i+1} = \omega$. □

Lemma 6. *Lemma 4 holds for all cyclic extensions E/F of degree 2 or 4.*

Proof. By the remark to Lemma 4, it suffices to prove that for E/F a cyclic extension of degree 4,

$$\rho^2 k_mE \cap (k_mE)^G \subseteq i_E N_{E/F} k_mE.$$

Let $\gamma \in K_mE$ such that $0 \neq \rho^2\bar{\gamma} \in \rho^2 k_mE \cap (k_mE)^G$. Then $l(\gamma) = 3$, and set $\beta := \rho\gamma$. We have

$$(\sigma^2 - 1)\bar{\beta} = \rho^3\bar{\gamma} = 0.$$

By Lemma 1 applied to E/E_1, $\bar{\beta} \in i_{E_1,E} k_mE_1$. Let $b \in K_mE_1$ such that $i_{E_1,E}\bar{b} = \bar{\beta}$.

Set $c := N_{E_1/F}b$. Then

$$i_{F,E}\bar{c} = i_{E_1,E}(\sigma + 1)\bar{b} = (\sigma^2 - 1)\bar{\gamma} = i_{E_1,E}N_{E/E_1}\bar{\gamma},$$

and since $l(\gamma) = 3$, $\langle \bar{\gamma} \rangle^G = \langle i_{F,E}\bar{c} \rangle$. By Lemma 1 we see that $i_{F,E_1}\bar{c} = N_{E/E_1}\bar{\gamma}$ and therefore $N_{E/E_1}\gamma = i_{F,E_1}c + 2g$ for some $g \in K_mE_1$.

Now $N_{E/F}\gamma = N_{E_1/F}(i_{F,E_1}c + 2g) = 2c + 2N_{E_1/F}g$. Set $\delta := b + g$. We calculate

$$N_{E/F}i_{E_1,E}\delta = 2c + 2N_{E_1/F}g,$$

so that $N_{E/F}(\gamma - i_{E_1,E}\delta) = 0$. By Theorem 1 there exists an $\alpha \in K_mE$ with $\rho\alpha = \gamma - i_{E_1,E}\delta$. Then, observing that $\rho^2(\overline{i_{E_1,E}\delta}) = 0$ since $\delta \in K_mE_1$, we see that

$$i_{F,E}N_{E/F}\bar{\alpha} = \rho^3\bar{\alpha} = \rho^2\overline{(\gamma - i_{E_1,E}\delta)} = \rho^2\bar{\gamma},$$

as required. □

3. Proof of Lemma 4

We prove Lemma 4 and hence, using the Reduction to Lemma 4, Theorem 2. We do so by induction on n, using Lemmas 5 and 6 as base cases and assuming the result for cyclic extensions of degree p^{n-1}. Observe that Lemmas 5 and 6 and the Reduction to Lemma 4 give us that Theorem 2 holds for degree p and, if $p = 2$, then for degree 4 as well. By the remark to Lemma 4, it suffices to prove that

$$\rho^{j-1} k_m E \cap (k_m E)^G \subseteq i_E N_{E/F} k_m E$$

for all $p^{n-1} < j < p^n$.

Assume that $k_m E = \oplus_{\gamma \in \Gamma} \mathbb{F}_p G\gamma$ is a decomposition of $k_m E$ into cyclic $\mathbb{F}_p G$-modules given by Proposition 1. Cyclic $\mathbb{F}_p G$-modules are indecomposable, and they are moreover self-dual and local, therefore with local endomorphism rings. By the Krull-Schmidt-Azumaya Theorem (see [AF, Theorem 12.6]), all decompositions of J into indecomposables are equivalent. (In our special case one can check this fact directly.)

First assume that $p > 2$. We know by Lemma 5 and by the Reduction to Lemma 4 that $k_m E|_{H_{n-1}}$ is a direct sum of trivial and free $\mathbb{F}_p H_{n-1}$-modules. Suppose that $\gamma \in k_m E$ satisfies $p^{n-1} < l(\gamma) < p^n$. We want to show that $\rho^{l(\gamma)-1}\, \gamma \in \rho^{p^{n-1}} k_m E = i_E k_m E$. Without loss of generality we may assume that γ generates a cyclic $\mathbb{F}_p G$-summand of $k_m E$. Since $l(\gamma) < p^n$, then by Lemma 3, $\mathbb{F}_p G\gamma|_{H_{n-1}}$ contributes direct summands which are not free or trivial, a contradiction. Hence a cyclic generator γ with length greater than p^{n-1} must have length p^n.

Now assume $p = 2$. Suppose that $\gamma \in \Gamma$ satisfies $2^{n-1} < l(\gamma) < 2^n$. By Lemma 6 and the Reduction to Lemma 4, $k_m E|_{H_{n-2}}$ is a direct sum of cyclic $\mathbb{F}_2 H_{n-2}$-modules of lengths 1, 2, and 4, and $\mathbb{F}_2 G\gamma$ is a cyclic summand of $k_m E$. By Lemma 3, we know that $\mathbb{F}_2 G\gamma|_{H_{n-2}}$ contributes cyclic $\mathbb{F}_2 H_{n-2}$-summands of length 3, a contradiction. Hence a cyclic generator $\gamma \in \Gamma$ of length greater than 2^{n-1} must have length 2^n.

In both cases, we have shown for $p^{n-1} < j \leq p^n$ that

$$\rho^{j-1} k_m E \cap (k_m E)^G = i_E N_{E/F} k_m E,$$

as desired. □

Acknowledgments

We are very grateful to A. Schultz as some of his ideas, developed in [MSS1], proved quite useful to us during the investigations leading to this paper. We also wish to thank M. Rost and J.-P. Serre for their comments after a lecture delivered in Nottingham by the third author in September 2005. These comments led to the improvement of our exposition.

References

[AF] F. Anderson and K. Fuller. *Rings and categories of modules.* Graduate Texts in Mathematics 13. New York: Springer-Verlag, 1973.

[BLMS] D. Benson, N. Lemire, J. Mináč, and J. Swallow. Detecting pro-p groups that are not absolute Galois groups. *J. Reine Angew. Math.*, to appear.

[BK] S. Bloch and K. Kato. p-adic étale cohomology. *Inst. Hautes Études Sci. Publ. Math.* **63** (1986), 107–152.

[Bo] Z. I. Borevič. The multiplicative group of cyclic p-extensions of a local field. *Trudy Mat. Inst. Steklov* **80** (1965), 16–29. Translated as *Proc. Steklov Inst. Math.* No. 80 (1965): Algebraic number theory and representations, D. K. Faddeev, ed., Providence, RI: American Mathematical Society (1968), 15–30.

[C] J. Carlson. *Modules and group algebras.* Lectures in Mathematics ETH Zürich. Basel: Birkhäuser Verlag, 1996.

[I1] O. Izhboldin. On the torsion subgroup of Milnor K-groups. *Dokl. Akad. Nauk SSSR* **294** (1987), no. 1, 30–33. Translated as *Soviet Math. Dokl.* **35**, no. 3, 493–495.

[I2] O. Izhboldin. On p-torsion in K_*^M for fields of characteristic p. *Algebraic K-theory.* Advances in Soviet Mathematics 4. Providence, RI: American Mathematical Society, 1991, 129–144.

[J] A. V. Jakovlev. Homological determination of p-adic representations of rings with basis of powers. *Izv. Akad. Nauk SSSR Ser. Mat.* **34** (1970), 1000–1014. Translated as *Math. USSR–Izvestiya* **4** (1970), no. 5, 1001–1016.

[L] T. Y. Lam. *Lectures on modules and rings.* Graduate Texts in Mathematics 189. New York: Springer-Verlag, 1999.

[LMS] N. Lemire, J. Mináč, and J. Swallow. Galois module structure of Galois cohomology and partial Euler-Poincaré characteristics. *J. Reine Angew. Math.*, to appear.

[Ma] B. Magurn. An algebraic introduction to K-theory. Encyclopedia of Mathematics and its Applications 87. Cambridge: Cambridge University Press, 2002.

[Mi] J. Milnor. Algebraic K-theory and quadratic forms. *Invent. Math.* **9** (1970), 318–344.

[MS] J. Mináč and J. Swallow. Galois module structure of pth-power classes of extensions of degree p. *Israel J. Math.* **138** (2003), 29–42.

[MSS1] J. Mináč, A. Schultz, and J. Swallow. Galois module structure of pth-power classes of cyclic extensions of degree p^n. *Proc. London Math. Soc.* **92** (2006), no. 2, 307–341.

[MSS2] J. Mináč, A. Schultz, and J. Swallow. Galois module structure of Milnor K-theory mod p^s in characteristic p. arXiv:math.NT/0602546v2.

[T] J. Thévenaz. Representations of finite groups in characteristic p^r. *J. Algebra* **72** (1981), 478–500.

Department of Mathematics, Physics and Engineering, Mount Royal College, 4825 Mount Royal Gate SW, Calgary, Alberta T3E 6K6 CANADA

E-mail address: `bhandari@mtroyal.ca`

Department of Mathematics, Middlesex College, University of Western Ontario, London, Ontario N6A 5B7 CANADA

E-mail address: `nlemire@uwo.ca`

E-mail address: `minac@uwo.ca`

Department of Mathematics, Davidson College, Box 7046, Davidson, North Carolina 28035-7046 USA

E-mail address: `joswallow@davidson.edu`